Divisible tilings in the Hyperbolic Plane

S. Allen Broughton
Rose-Hulman Institute of Technology


Divisible Tilings

- Divisible euclidean tilings
- hyperbolic geometry, area of a polygon, tilings
- divisible tilings, formula for K
- free and bound vertices, special K
- combinatorial search
- geometric search
- further questions

Divisible euclidean tilings

- show examples
- show tiling on a surface (further question)

(2,4,4) -tiling of the torus



Kaleidoscopic Tiling of the Plane: Definition by example

- Tiling: Covering by polygons "without gaps and overlaps"
- Kaleidoscopic: Symmetric via reflections in edges.
- Geodesic edges extend to lines in the tiling
- Kaleidoscopic polygons if and only angles of the form π where n is an integer

n

Kaleidoscopic Tiling of the Plane: Terminology

- terminology
 - (l,m,n) -triangle,
 - (s,t,u.v)-quadrilateral

Hyperbolic geometry

- Points, lines and angles
- reflections show picture
- formula for area of a triangle

$$\pi - \left(\frac{\pi}{l} + \frac{\pi}{m} + \frac{\pi}{n}\right)$$

$$= \pi \left(1 - \frac{1}{l} - \frac{1}{m} - \frac{1}{n}\right)$$

Hyperbolic geometry

formula for area of a quadrilateral

$$2\pi - \left(\frac{\pi}{s} + \frac{\pi}{t} + \frac{\pi}{u} + \frac{\pi}{v}\right)$$

$$= \pi \left(2 - \frac{1}{s} - \frac{1}{t} - \frac{1}{u} - \frac{1}{v}\right)$$

Tilings and divisible tilings - 1

- (2,3,7) and (3,3,4) example of tilings
- show divisible tilings created from (2,3,7)-example
- Divisible tiling if tiling can be kaleidoscopically subdivided by a finer tiling

Tilings and divisible tilings - 2

 Divisible tilings can be found by kaleidoscopically subdividing a kaleidoscopic tile by another kaleidoscopic tile

Formula for K

• K = number of tiles required to tile a larger tile

$$K = \frac{\pi(2 - \frac{1}{l} - \frac{1}{l} - \frac{1}{l})}{s t u v}$$

$$\frac{\kappa}{\pi(1 - \frac{1}{l} - \frac{1}{l} - \frac{1}{l})}$$

Free and bound vertices, Special K

- □ Free vertices, infinite families of tilings, show example
- bound vertices, finitely many examples
- □ If K > special K there are only bound vertices
- □ Special K = 12

Combinatorial search K <= special K

- Show associated Catalan and hub polygons
- □ work out K=4, combinatorially
- C(12) = 208012 so there are that many Catalan polygons
- use dihedral symmetry to reduce to 7528

Geometric/Algebraic search K > special K

- \square show (2,3,7) example for (7,7,7)
- \square show failed (2,3,7) tiling of (7,7,7,7)
- \square show algebraically why (2,3,7) tiles (7,7,7)